Dissecting an Amphibian—the Frog

50

BACKGROUND

Frogs are members of the class Amphibia. Biologists have appropriately named these organisms amphibians because they have adaptations that enable them to "live in two ways"—both on land and in water. In this laboratory you will examine a frog and observe how it is adapted to both environments.

OBJECTIVES

- Dissect and identify the external and internal structures of a frog.
- List three ways the frog is adapted to life on land and in water.

MATERIALS

Preserved frog

Hand lens

Dissecting tray

Compound microscope

Dissecting pins

Stereoscopic dissecting microscope

Dissecting scissors

Probe

Forceps

Dissecting needle


PROCEDURE A

THE EXTERNAL ANATOMY OF THE FROG

- 1. Place the preserved frog in your dissecting tray. Obtain all the dissecting equipment and place it at your work area.
- 2. Observe the frog. Feel the smooth and pliable skin. Note the coloration on all parts of the body. Describe how the coloration on the dorsal surface and ventral surface differ.

(a)


3. The division of a frog's body includes the head, trunk, and limbs or legs as shown in a. All frogs lack necks and tails. Examine the front and hind limbs of the frog. Find the digits which are fingerlike projections on the forelimbs and hindlimbs. Locate the hindlimbs which are the longer and more muscular limbs of the frog. A hindlimb consists of a thigh, shank, ankle, foot, and webbed toes. Locate the forelimbs which are shorter than the hindlimbs. A forelimb consists of an upper arm, wrist, hand, and fingers. Find on the male frog the male pad located on the innermost finger of the forelimb. The male pad is a dark swelling or enlargement used by a male to grasp a female during mating.




4. Examine the frog's head. Notice the size of the frog's eyes and how the eyes protrude. Find the nonmoveable upper and lower eyelids. Locate the nictitating membrane that is a transparent covering that sweeps upward over the eyes. Identify the tympanic membrane or the eardrum located on each side of the head. Observe the frontal organ or brow spot found between and anterior to the two eyes. The frontal organ is the remaining indication of a third or middle eye. Note the external nares or nostrils that are the openings on the anterior and dorsal part of the skull. Label each of these structures on **b**: eyes, nictitating membrane, tympanic membrane, frontal organ, external nares.

PROCEDURE B

OBSERVING THE FROG'S MOUTH CAVITY





c. The frog's mouth cavity

1.	Pry open the mouth. Use the scissors to cut the
	corners of the mouth where the maxilla (upper
	jaw) and mandible (lower jaw) join together. Ex-
	amine the frog's mouth. Locate the tongue which
	is a muscular, sticky flap on a living frog. The
	frog flicks out the tongue from the floor of the
	mouth to catch flying insects. Flip out the tongue.
	Describe how the tongue is attached to the
	mouth.

(b)				

Feel the *maxillary teeth* that are found along the rim of the upper jaw or *maxilla*. Describe how the maxillary teeth would be useful to the frog.

/ \			
(C)			

Find the *vomerine teeth* attached to the skull bones of the roof of the mouth. Notice that only the upper jaw has small teeth. Locate the *internal nares* or *nostrils* found in the roof of the mouth. How is the location of the nares an adaptation to living in water?


(d)			
(α)			

Locate the glottis which is a slitlike opening on the swollen voice box found on the floor of the mouth. Air passes through the glottis going to and from the lungs. Find the esophagus at the rear of the mouth. The wide opening of the esophagus allows the frog to swallow food whole. The esophagus is the first tube of the alimentary canal leading to the stomach. Notice each eustachian tube opening found lateral to the esophagus near the hinge of the upper jaw. The eustachian tube openings lead to each ear and ensure equal air pressure on both sides of the tympanic membrane. Label each of these structures on c: tongue, maxillary teeth, vomerine teeth, internal nare, glottis, esophagus, and eustachian tube opening.

50

Dissecting an Amphibian— The Frog (Continued)

PROCEDURE C



d. How to dissect the frog

THE INTERNAL ANATOMY OF THE FROG

Place the frog in the dissection tray ventral side up and pin down both jaws.
 Use the forceps and scissors to lift a piece of skin where the hind legs meet. Insert the scissors and cut just the skin along the midline to the level of the lower jaw as shown in d. Clip the skin at a right angle to the incision as shown. Observe the layers of muscles now exposed. Repeat the incision procedure for the muscle tissue. Make this incision a little to the right of center to avoid cutting a major vein.


(NOTE: Always lift the parts to be cut so as to avoid damaging underlying tissues and organs.) In the chest region you will have to cut through the breast-bone. Again make right angle cuts from each end of the long incision as shown in **d.** Fold open and pin down flaps of the skin and body wall. Make additional cuts in the breastbone, if necessary, to expose the organs in the upper body.

2. The sex of the frog is readily noticed once the internal organs are exposed. If there is a mass of black and yellow eggs in the transparent ovaries, the frog is a female. If the ovaries obscure the view of the other organs the ovaries may be cut out of the body at their point of attachment and removed from the body cavity.

Respiratory System

3. Locate the small pair of lungs and the bronchial tubes as shown in **e.** Describe the texture of the lungs.

(e) _____ Label the lung as shown in **e.**




Circulatory System

4. Find the heart in the center of the chest cavity between the lungs. Notice that the heart lies in a thin sac called the pericardium. Remove the pericardium to observe the three-chambered heart. The two, dark-walled chambers called atria receive blood. The right atrium receives deoxygenated blood from the veins of the frog's body. The left atrium receives oxygenated blood from the lungs. Each atrium empties into the ventricle which is the lightercolored, thick-walled part of the heart. The large vessel forming a y-shape at the anterior end of the ventricle is the conus arteriosus. Blood is pumped out of the conus arteriosus through a system of arteries that you see around the heart. Label the right atrium, left atrium, ventricle, and conus arteriosus on e.

Digestive System

5. Find the dark brown liver composed of three to five lobes. Between the right lobe and the one next to it, is the transparent gall bladder, usually filled with the green bile that it stores. Locate the bile duct that brings bile from the gall bladder to the first part of the small intestine or duodenum. The duodenum and stomach form a loop that is held in place by sheets of membranous connecting mesentery. In the mesentery, between the stomach and duodenum, lies the pancreas, a glandular organ that produces digestive enzymes. The pancreas empties its enzymes through a duct that empties into the bile duct. The dark, spherical spleen, which is a circulatory and lymphatic system organ, lies in the mesentery posterior to the stomach. One of the spleen's functions is to store blood. The small intestine proceeds posteriorly to the coiled ileum that widens into the large intestine. Where the urinary bladder and reproductive ducts empty into the intestine, the section is called the cloaca. The wastes in the cloaca empty out at the anus. Label each of the following structures in f: liver, gall bladder, bile duct, duodenum, ileum, small intestine, stomach, mesentery, pancreas, large intestine, cloaca, anus, spleen, and urinary bladder.




f. The internal anatomy of the frog

Dissecting an Amphibian— The Frog (Continued)

50

Excretory System

6. Push the intestines aside to see the two dark, oblong kidneys lying on the dorsal wall of the body cavity. Find the ureter, which is a tube leading out of each kidney. The ureters carry the chemical wastes from the blood that the kidneys have filtered out to the urinary bladder. The chemical wastes called urine are stored in the urinary bladder that empties into the cloaca. The adrenal gland, which is part of the endocrine system, can be seen as a yellow stripe on each kidney. Fat bodies, whose purpose is to store fat to be used during hibernation or when food is scarce, attach to the anterior end of each kidney. Usually fat bodies are bright yellow, finger-like projections. Label the kidneys, ureters, cloaca, adrenal glands, and fat bodies in g.



g. The reproductive and excretory systems of a male and a female frog

Reproductive System

- 7. In male frogs each yellowish, oval-shaped *testis* is attached by tubes to the kidney. During mating the sperm mixes with and follows the exit route of the urine. The males of many frog species have vestigial oviducts attached to the cloaca called *Mullerian tubes*. In female frogs the two ovaries appear as *egg* sacs. Eggs released from the ovaries enter a long, coiled *oviduct* at the anterior opening. Eggs pass down the cream-colored oviduct and are held in an *ovisac* or *uterus* before entering the cloaca. From the cloaca the eggs pass from the body to be fertilized outside the frog's body. Label the *Mullerian tube*, *egg* sac, *oviduct*, *ovisac*, and *testis* on **g**.
- 8. Dispose of your dissected frog where your teacher instructs. Clean up your work area and equipment. Return the cleaned dissecting equipment to the appropriate place.

ANALYSIS	1.	List the frog's features that are similar to features of animals that live in water.
	2.	How might the countershading of a frog help protect it from predators?
	3.	List the organs of the frog's digestive system through which swallowed food passes.
	4.	Which glands are part of the frog's digestive system?
		Trace the path of urine through the frog's body beginning with the organ that filters waste from the blood.
	6	List three ways the frog is adapted to life on land.

GOING FURTHER

- 1. Observe a living tadpole. Find out how the structures that a tadpole uses for locomotion, respiration, and ingestion of food differ from those of a frog.
- 2. Remove the brain of the frog and make a sketch of the dissected brain. Use a reference book to identify the parts of the brain.

